## NICHOLAS BASTIDAS, MD, FAAP

Plastic, Reconstructive & Craniofacial Surgery Associate Professor of Surgery Manhattan Office 210 East 68th St., Suite 1F New York, NY 10065 (T) 212.485.0616 Long Island Office 1991 Marcus Ave., Suite 102 Lake Success, NY 11042 (T) 516-497-7900 (F)516.497.7920

## What to expect after Facial Feminization Procedures:

- You will probably spend 1- 2 days in the hospital after surgery. We will attempt to get you a
  private room.
- You can expect to be uncomfortable for 7-10 days. You will be given a prescription for pain medication, most likely oxycodone with acetaminophen. You can take 1-2 pills every 4-6 hours. Do not drive while taking prescription pain medication. After 48 hours, you can also take Advil or Ibuprofen.
- Take all medicine with food. Pain medication can cause constipation. You can take Miralax or Dulcolax (over the counter) if needed.
- You can shower normally when you go home and let soap and water run over any incision sites.
- Apply a small amount of bacitracin or Neosporin twice daily to any incision site on your head or scalp if there are no surgical tapes covering the incisions.
- Sleep with your head elevated on at least 2-3 pillow for the first 3 nights. Do not lay flat for naps during the day. Do not lay on your stomach with your face down.
- If you have stitches in your mouth, rinse with Peridex (prescription mouthwash) twice daily or dilute Listerine (Mix ½ Listerine with ½ water) for 7 days.
- If you have stitches in your mouth, eat a mainly liquid or very, very soft diet for at least 1 week.
- If you have had any type of chin or jaw surgery, do not eat any hard foods, or anything that requires a lot of chewing for 2 weeks.
- Gently brush teeth twice daily and rinse mouth with water every time after eating or drinking.
- No bending or heavy lifting. Keep your head upright and above your heart. No strenuous exercise.
- You can expect bruising around your eyes, mouth and jaw. Apply ice for 20 minutes every 1-2 hours for the first two days after surgery.
- Leave the jaw bra (chin/face support garment) in place except briefly to shower for 2 weeks.

## Please call the office or Nurse Practitioner (347-502-1439) if...

- You have a fever greater than 100.5 degrees F
- You have continuous bleeding that won't stop with ice or pressure
- You notice a funny odor, or yellow, pus-like discharge
- There is redness, warmth or opening on any incision sites
- If are unable to tolerate any fluids and have persistent vomiting

## IN CASE OF AN EMERGENCY OVERNIGHT, THERE IS ALWAYS A DOCTOR ON CALL THAT CAN BE REACHED BY CALLING THE MAIN OFFICE NUMBER (516-497-7900)